

MYCOTAXON

<http://dx.doi.org/10.5248/123.265>

Volume 123, pp. 265–270

January–March 2013

***Peltaster fructicola*, a newly recorded species from China associated with sooty blotch and flyspeck**

CHEN CHEN¹, LIU GAO¹, MEIYAN QU¹, XIAOYONG WEI¹,
WENHUAN LI¹, RONG ZHANG¹, GUANGYU SUN^{1*} & MARK L. GLEASON²

¹State Key Laboratory of Crop Stress Biology in Arid Areas and College of Plant Protection,
Northwest A&F University, Yangling, Shaanxi 712100, P. R. China

²Department of Plant Pathology and Microbiology, Iowa State University,
Ames, Iowa 50011, U.S.A

* CORRESPONDENCE TO: sgy@nwsuaf.edu.cn

ABSTRACT — *Peltaster fructicola* is first reported and described from China. This fungus was isolated from peels of crabapple and hawthorn collected from Liaoning and Shaanxi Provinces and exhibited the punctate mycelial type of sooty blotch and flyspeck on both hosts. Our isolates were identified based on morphological characteristics and DNA sequence analysis.

KEY WORDS — *Rosaceae*, taxonomy, morphology, phylogeny

Introduction

Sooty blotch and flyspeck (SBFS) is a commonly occurring disease complex in humid and temperate regions worldwide, causing blemishes on apple, pear, banana, and other cultivated crops that result in economic losses (Williamson & Sutton 2000, Batzer et al. 2005, Gleason et al. 2011). The SBFS fungi colonize the waxy cuticle of plants and display various mycelial types. The term “sooty blotch” refers to colonies on hosts that consist of a dark mycelial mat with or without sclerotium-like bodies, whereas “flyspeck” denotes a cluster of black dots without a visible mycelial mat. “Sooty blotch” was attributed to a single fungal species for more than 75 years (Colby 1920, Williamson & Sutton 2000) until the late 1990s, when Johnson et al. (1996, 1997) found three “sooty blotch” species based on morphological examination of sooty blotch isolates from apples in USA. One newly described species from that work was *Peltaster fructicola* (Johnson et al. 1996).

During a recent survey of SBFS complex on fruit of rosaceous hosts in China, crabapples and hawthorns were found to display colonies of SBFS. Based on analysis of morphology and DNA sequences of nuclear ribosomal ITS region, our isolates were identified as *P. fructicola*.

Materials & methods

Isolates and morphology

Crabapple (*Malus ×micromalus*) and hawthorn (*Crataegus pinnatifida*) fruit with SBFS signs were collected from Liaoning and Shaanxi Provinces. Two isolates were obtained from hawthorns in Liaoning and Shaanxi Provinces, and three from crabapples in Liaoning Province. Peels displaying SBFS were examined, photographed and preserved. Thalli were transferred directly from peels to potato-dextrose agar (PDA) and cultured at 25°C in the dark (Sun et al. 2003). After one month, pure isolates were transferred to PDA and water agar (WA, 2% agar) for morphological examination. Colony descriptions were based on cultures on PDA after 3 weeks in the dark at 25°C. Morphological descriptions were made on WA and synthetic nutrient-poor agar (SNA) after 3 weeks using the method of Zhang et al. (2009). Specimens and representative dried culture were deposited in the Fungal Herbarium of Northwest A&F University (HMUABO), Yangling, Shaanxi Province, China.

DNA extraction, PCR and sequencing

Genomic DNA was extracted from fungal mycelium according to the protocol of Barnes et al. (2001). The primers ITS1-F (Gardes & Bruns 1993) and ITS4 (White et al. 1990) were used to amplify the internal transcribed spacer (ITS) region of nuclear ribosomal DNA. The PCR conditions followed the methods of Zhuang et al. (2010). PCR products were sequenced by Sangon Biotech (Shanghai), China.

Sequence alignment and phylogenetic analysis

Sequences generated in this study were added to other sequences of *Peltaster* spp., as well as the outgroup species *Mycosphaerella lateralis* and *Schizothyrium pomi* obtained from GenBank. Preliminary alignments were performed using CLUSTAL X (Thompson et al. 1997), and then manually adjusted using BioEdit v. 5.0.9.1 (Hall 1999). Phylogenetic analysis of aligned sequences was performed using PAUP v. 4.0b 10 (Swofford 2003). The heuristic search option was 1000 random taxa addition and tree bisection-reconnection (TBR) as the branch-swapping algorithm. The robustness of clades and internal branches was evaluated by 1000 bootstrap replications. Tree length (TL), consistency index (CI), retention index (RI) and rescaled consistency index (RC) were also calculated. Sequences generated in this study were deposited in GenBank. Alignment and the representative tree were deposited in TreeBase (<http://purl.org/phylo/treebase/phyloids/study/TB2:S13540>).

Results

Phylogenetic analysis

The manually adjusted ITS alignment contained 27 sequences (including the outgroup sequences). Of the 544 characters used for the phylogenetic analysis, 153 were parsimony-informative, 95 were variable and parsimony-uninformative, and 296 were constant. The first 100 equally most parsimonious trees were obtained from the parsimonious analysis, and the first of which is shown in FIG. 1 (TL = 403 steps, CI = 0.8734, RI = 0.9373, RC = 0.8187). The *Peltaster* species formed a strongly supported clade with a bootstrap value of




FIG. 1. One of 100 equally most parsimonious trees obtained from a heuristic search with 1000 random taxon additions of the ITS sequence alignment. The bootstrap support values (50%) based on 1000 replicates are shown at the nodes. The tree was rooted to *Mycosphaerella lateralis* and *Schizothyrium pomi*. The scale bar shows 10 changes. Strains investigated in this paper are presented in bold.

100%, which further divided into two subclades. Our five strains clustered together with *Peltaster fructicola* within one of the subclades with a high bootstrap value of 100%, indicating that they might represent the same species (FIG. 1).

Taxonomy

Peltaster fructicola Eric M. Johnson, T.B. Sutton & Hodges,

Mycologia 88(1): 120, 1996

FIG. 2

HYPHAE smooth, septate, branched, hyaline and turning to pale brown as culture aged, 1.5–2.0 µm wide. CONIDIOPHORES absent. CONIDIOGENOUS CELLS hyaline, monoblastic, intercalary, formed directly on undifferentiated hyphae. Conidiogenous loci inconspicuous, stub-shaped, concolorous with conidiogenous cells. CONIDIA solitary, unicellular, hyaline, elliptic to ovoidal, produced in basipetal order from the conidiogenous loci and aggregated in

masses near the conidiogenous cells, $4\text{--}6.5(-7) \times 1.5\text{--}2.5\text{ }\mu\text{m}$. Microcyclic conidiation observed in culture, with expansion and germination of conidia. Conidia on WA slightly smaller in size than those on SNA, $4\text{--}6(-7) \times 1.5\text{--}2.0\text{ }\mu\text{m}$.

CULTURAL CHARACTERISTICS — On PDA erumpent, compact, with smooth, crenate margin and sparse to no aerial mycelium, surface strongly folded, outer zone light gray, center greenish gray to olivaceous-gray, reverse black with yellow masses of conidia at the bottom of the colony; reaching (13–)14(–16) mm diam after 21 d at 25°C in the dark.

On peels of crabapple fruit, showing punctate mycelial type, visible mycelial mat with dull brown, flattened, circular or irregular sclerotium-like bodies, 68–145 μm diam. On peels of hawthorn fruit similar, sclerotium-like bodies 62–223 μm diam.

SPECIMENS EXAMINED (all conserved in HMUABO): **CHINA, LIAONING PROVINCE:** Huludao City, Suizhong County, $40^{\circ}19'32''\text{N}$ $120^{\circ}17'28''\text{E}$, on fruit surface of crabapple (*Malus $\times$ micromalus* Makino), Sept. 2010, Zhuang JL LNHT1506 (GenBank JX961608), LNHT1802 (GenBank JX961609), LNHT1405a (GenBank JX961610); on fruit surface of hawthorn (*Crataegus pinnatifida* Bunge), Sept. 2010, Zhuang JL LNSZ1301a (GenBank JX961607). **SHAANXI PROVINCE:** Shangluo City, Shangnan County, on fruit surface of hawthorn (*Crataegus pinnatifida*), 21 Oct. 2011, Chen C & Dang JL SNSZ17 (GenBank JX961611).

Discussion

The genus *Peltaster* was established by Sydow & Sydow in 1917; the type species is *Peltaster hedytidis* Syd. & P. Syd. Sutton and his colleagues described *P. fructicola* (Johnson et al. 1996) as one of the pathogens of sooty blotch and later associated it with punctate mycelial type on apple (Williamson et al. 2004). Since then, this fungus has been found widely in the eastern and midwestern United States (Johnson et al. 1997, Batzer et al. 2005, Díaz Arias et al. 2010). As the etiology of SBFS is studied worldwide, *P. fructicola* has been reported to cause sooty blotch in many other countries and regions, including Serbia, Montenegro, and Poland (Ivanović et al. 2010, Mirzwa-Mróz & Wińska-Krysiak 2011), indicating that *P. fructicola* is among the most prevalent SBFS species.

Based on phylogenetic analysis of the ITS region and morphological comparison, we identified our isolates as *Peltaster fructicola*, a newly recorded species for China. To our knowledge, this is also the first report of SBFS on crabapple and hawthorn caused by *P. fructicola*.

Acknowledgments

This work was supported by National Natural Science Foundation of China (31170015, 31171797), the 111 Project from Education Ministry of China (B07049), Top Talent Project of Northwest A&F University and the earmarked fund for Modern Agro-industry Technology Research System (nycytx-08-04-04). The authors wish to thank Dr Eric H.C. McKenzie (Landcare Research, Private Bag 92170, Auckland,




FIG. 2. *Peltaster fructicola* (LNHT1506). A. Signs on crabapple. B. Colony on PDA. C, D. Conidia and conidiogenous cells. E–G. Conidiogenous cells giving rise to conidia, with visible conidiogenous loci. H–J. Conidia expanding and germinating and microcyclic conidiation. Scale bars: A = 0.2 mm; C–E, H–J = 10 μ m; F, G = 5 μ m.

New Zealand) and Professor Zhongyi Zhang (College of Plant Protection, Yunnan Agricultural University, Kunming, Yunnan, China) for reviewing the manuscript.

Literature cited

- Barnes I, Roux J, Wingfield MJ, Coetzee MPA, Wingfield BD. 2001. Characterization of *Seiridium* spp. associated with cypress canker based on β -tubulin and histone sequences. *Plant Disease* 85: 317–321. <http://dx.doi.org/10.1094/PDIS.2001.85.3.317>
- Batzer JC, Gleason ML, Harrington TC, Tiffany LH. 2005. Expansion of the sooty blotch and flyspeck complex on apples based on analysis of ribosomal DNA gene sequences and morphology. *Mycologia* 97: 1268–1286. <http://dx.doi.org/10.3852/mycologia.97.6.1268>

- Colby AS. 1920. Sooty blotch of pomaceous fruits. Transactions of the Illinois State Academy of Science 13: 139–175.
- Díaz Arias MM, Batzer JC, Harrington TC, Wong AW, Bost SC, Cooley DR, Ellis MA, Hartman JR, Rosenberger DA, Sundin GW, Sutton TB, Travis JW, Wheeler MJ, Yoder KS, Gleason ML. 2010. Diversity and biogeography of sooty blotch and flyspeck fungi on apple in the eastern and midwestern United States. *Phytopathology* 100: 345–355. <http://dx.doi.org/10.1094/PHYTO-100-4-0345>
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2: 113–118. <http://dx.doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Gleason ML, Batzer JC, Sun GY, Zhang R, Díaz Arias MM, Sutton TB, Crous PW, Ivanović M, McManus PS, Cooley DR, Mayr U, Weber RWS, Yoder KS, Del Ponte EM, Biggs AR, Oertel B. 2011. A new view of sooty blotch and flyspeck. *Plant Disease* 95: 368–383. <http://dx.doi.org/10.1094/PDIS-08-10-0590>
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Ivanović MM, Ivanović SM, Batzer JC, Tatalović N, Oertel B, Latinović J, Latinović N, Gleason ML. 2010. Fungi in the apple sooty blotch and flyspeck complex from Serbia and Montenegro. *Journal of Plant Pathology* 92: 65–72.
- Johnson EM, Sutton TB, Hodges CS. 1996. *Peltaster fruticola*: a new species in the complex of fungi causing apple sooty blotch disease. *Mycologia* 88: 114–120. <http://dx.doi.org/10.2307/3760790>
- Johnson EM, Sutton TB, Hodges CS. 1997. Etiology of apple sooty blotch disease in North Carolina. *Phytopathology* 87: 88–95. <http://dx.doi.org/10.1094/PHYTO.1997.87.1.88>
- Mirzwa-Mróz E, Wińska-Krysiak M. 2011. Diversity of sooty blotch fungi in Poland. *Acta Scientiarum Polonorum-Hortorum Cultus* 10: 191–200.
- Sun GY, Zhang R, Zhang Z, Zhang M. 2003. Isolation of sooty blotch and flyspeck fungi from apple surface by picking up the thalli. *Acta Phytopathologica Sinica* 33: 479–480.
- Swofford DL. 2003. PAUP*. Phylogenetic analysis using parsimony (*and other methods). Version 4.0. Sinauer Associates, Sunderland, Massachusetts, USA.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25: 4876–4882. <http://dx.doi.org/10.1093/nar/25.24.4876>
- White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis et al. (eds.), *PCR Protocols: a guide to methods and applications*. San Diego, Academic Press.
- Williamson SM, Sutton TB. 2000. Sooty blotch and flyspeck of apple: etiology, biology, and control. *Plant Disease* 84: 714–724. <http://dx.doi.org/10.1094/PDIS.2000.84.7.714>
- Williamson SM, Hodges CS, Sutton TB. 2004. Re-examination of *Peltaster fruticola*, a member of the apple sooty blotch complex. *Mycologia* 96: 885–890. <http://dx.doi.org/10.2307/3762121>
- Zhang R, Yang HL, Sun GY, Li HY, Zhuang JL, Zhai XR, Gleason ML. 2009. *Strelitziana mali*, a new species causing sooty blotch on apple fruit. *Mycotaxon* 110: 477–485. <http://dx.doi.org/10.5248/110.477>
- Zhuang JL, Zhu MQ, Zhang R, Yin H, Lei YP, Sun GY, Gleason ML. 2010. *Phialophora sessilis*, a species causing flyspeck signs on bamboo in China. *Mycotaxon* 113: 405–413. <http://dx.doi.org/10.5248/113.405>